



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

REMARKS BY TRACY A. PIERCE, Harvard University.

It is well known that every prime of the form $4m + 1$ is the sum of two squares. Using Lucas's form of an odd perfect number (see Problem 211 above) we see that

$$(4m + 1)^{4k+1}n^2 = (x^2 + y^2)[(4m + 1)^{2k}n^2 = (x^2 + y^2)N^2 = X^2 + Y^2,$$

contrary to the proposition as proposed.

229. (March, 1915.) Proposed by WALTER C. EELLS, Whitman College.

If p and q are integers and p is prime and positive, find the condition on q that the equation $p^x = qx$ shall have integral solutions, solve for x , and show that for a special value of p it has two solutions for a certain q , otherwise only one.

I. SOLUTION BY FRANK IRWIN, University of California.

Since p is prime, we must have $x = p^a$, $q = p^b$, where a and b are positive integers or zero (for a). Now $p^{a+b} = qx = p^{p^a}$, so that $b = p^a - a$, and q is necessarily of the form $p^{(p^a - a)}$. This condition is evidently also sufficient.

Given, then, such a q , the exponent $p^a - a$ may be determined; then a , which will give us x , is that number which must be added to this exponent to make it equal to the *next higher* power of p . For no power of p can lie between $p^a - a$ and p^a , since $p^a - a > p^{a-1}$, as may be readily proved, for instance by mathematical induction.

Of the cases that require special investigation, $a = 0$, and $p = 2$ with $a = 1$ or 2 , the only one for which, given $p^a - a$, there is more than one solution for a , is the case

$$2^a - a = 1,$$

which has two solutions $a = 0, 1$. There are two solutions of our problem then for the case $p = 2$, $q = 2$, viz., $x = 1, 2$.

II. SOLUTION BY THE PROPOSER.

Consider the two functions, $y = p^x$, $y = qx$. For $x = k$ (any integer), $y = p^k$. The slope of line $y = qx$, passing through (k, p^k) , is $q = p^k/k$, which is integral if and only if

$$k = p^n \quad (n = 0, 1, 2, 3, \dots), \text{ i. e., } q = p^{(p^n - n)}.$$

(If $k < 0$, q is fractional since it is then $= 1/kp^k$).

Substituting this value of q in the given equation, it is easily seen that it is satisfied if and only if

$$x = p^n \quad (n = 0, 1, 2, 3, \dots).$$

Consider the exponent of p , namely $p^n - n$. We have

$$[p^n - n]_{n=0} = 1, \text{ and } [p^n - n]_{n=1} = p - 1.$$

Then $x_1 = p^0$ and $x_2 = p^1$ will be solutions of the given equation if $1 = p - 1$, i. e., if $p = 2$. From the graphs of the exponential function it is easily seen that $y = qx$ can have but one integral intersection if $p \neq 2$, $n_1 \neq 0$, $n_2 \neq 1$.

The equation having two solutions is $2^x = 2x$, of which $x_1 = 1$, $x_2 = 2$.

230. (April, 1915.) Proposed by E. B. ESCOTT, Ann Arbor, Michigan.

Find three numbers such that their sum, the sum of their squares, and the sum of their cubes, shall be a cube.

Note.—W. D. Cairns says this problem, which was proposed in *L'Intermédiaire* in 1900, remains unsolved to date, even though it was reprinted in February, 1913.

REMARKS BY ARTEMAS MARTIN, Washington, D. C.

The above problem was published in the *Mathematical Visitor*, Vol. I, No. 1 (Erie, Pa., March, 1877), page 6, as No. 9 in a list of "Unsolved Problems." So far as the writer at present knows that was the first publication of the problem and it still remains "unsolved."